

Species *Methanocatella smithii*^{Ts}

Etymology

[smith'i.i] N.L. gen. masc. n. *smithii*, of Smith, named after P.H. Smith, who isolated the type strain

Nomenclatural type

[NCBI Assembly: GCF_000016525.1](#)^{Ts}

Reference Strain

[ATCC 35061](#) = PS = [DSM 861](#)

Description

The species identified by metagenomic analyses. The G+C content of the type genome is 31.0 mol%, and the genome size is 1.85 Mbp. Cells are short oval rods or coccobacilli with tapered ends, 0.6–0.7 µm in width and ~1.0 µm in length. Cells occur most frequently in pairs or in chains of 4–6 cells. Gram positive. Nonmotile. H₂ and CO₂ are the preferred energy sources. Growth on formate is poor. Cells grow optimally at 37 to 39 °C. (as in [Balch et al., 1979](#); [Miller, 2015](#)).

Classification

Archaea » *Methanobacteriota* » *Methanobacteria* » *Methanobacteriales* » *Methanobacteriaceae* » *Methanocatella* » *Methanocatella smithii*^{Ts}

References

Effective publication: Protasov et al., 2023 [1]

Registry URL

<https://seqco.de/i:32433>

References

1. Protasov et al. (2023). Diversity and taxonomic revision of methanogens and other archaea in the intestinal tract of terrestrial arthropods. *Frontiers in Microbiology*. [DOI:10.3389/fmicb.2023.1281628](https://doi.org/10.3389/fmicb.2023.1281628)