

Species *Methanorbis furvi*^{TS}

Etymology

[fur'vi] L. gen. masc. n. *furvi*, of the black one, referring to the color of the host.

Nomenclatural type

[NCBI Assembly: GCA_032714615.1](#)^{TS}

Reference Strain

[Strain sc|0038661](#): Ag1 = [DSM 115764](#)

Description

Ovoid cocci 1–3 µm. Non-motile. H₂ and CO₂ are the substrates for methanogenesis; formate can substitute H₂. Acetate and formate are required for growth. The optimum temperature is 33 °C. The G+C content of the type genome is 50.1 mol%, and the genome size is 1.84 Mbp.

Classification

Archaea » *Methanobacteriota* » “Methanomicrobia” » *Methanomicrobiales* » *Methanocorpusculaceae* » *Methanorbis* » *Methanorbis furvi*^{TS}

References

Effective publication: Protasov et al., 2023 [1]

Registry URL

<https://seqco.de/i:32572>

References

1. Protasov et al. (2023). Diversity and taxonomic revision of methanogens and other archaea in the intestinal tract of terrestrial arthropods. *Frontiers in Microbiology*.
[DOI:10.3389/fmicb.2023.1281628](https://doi.org/10.3389/fmicb.2023.1281628)