

Species *Methanolapillus millepedarum*^{Ts}

Etymology

[mil.le.pe.da'rum] L. gen. pl. n. *millepedarum*, of millipedes

Nomenclatural type

[NCBI Assembly: GCA_032594115.1](#)^{Ts}

Reference Strain

Ac7

Description

Irregular cocci cells 1–2 µm in diameter, non-motile. Use methanol and methylamines as methanogenesis substrates together with hydrogen.). Requires acetate, yeast extract, Casamino acids, and coenzyme M. The optimum temperature is 37 °C. The G+C content of the type genome is 42.9%, and the estimated genome size is 1.93 Mbp.

Classification

Archaea » *Methanobacteriota* » “Methanomicrobia” » *Methanosarcinales* » *Methanosarcinaceae* » *Methanolapillus* » *Methanolapillus millepedarum*^{Ts}

References

Effective publication: Protasov et al., 2023 [1]

Registry URL

<https://seqco.de/i:32600>

References

1. Protasov et al. (2023). Diversity and taxonomic revision of methanogens and other archaea in the intestinal tract of terrestrial arthropods. *Frontiers in Microbiology*. [DOI:10.3389/fmicb.2023.1281628](https://doi.org/10.3389/fmicb.2023.1281628)