

---

## Species *Candidatus* *Adiutrix* intracellularis

---

### Etymology

L. *intra*, within; L. *fem. n. cellula*, a small chamber or cell; L. *fem. suff. -aris*, suffix denoting pertaining to; N.L. *fem. adj. intracellularis*, intracellular

### Nomenclatural type

Unknown

### Description

Properties: Rod-shaped bacteria (approximately 0.5–0.6  $\mu$ m in diameter and 0.8–1.9  $\mu$ m in length) with slightly pointed ends. Form a monophyletic group with the SSU rRNA genes of other *Deltaproteobacteria* from termite guts. Possesses genes coding for production of acetate from CO<sub>2</sub> and H<sub>2</sub> (Wood–Ljungdahl pathway) and nitrogen fixation. Colonize the cytoplasm of the parabasalid flagellate *Trichonympha collaris* in the hindgut of the termite *Zootermopsis nevadensis*.

So far uncultured. The basis of assignment is the SSU rRNA gene sequences of representative phylotypes (Accession No. AB972401; AB894435–AB894480) and hybridization with the specific SSU rRNA-targeted oligonucleotide probe Delta-Tr3-Zn (5'-CTT GAA CCG AAG TTC CTG-3'). A draft genome of strain Adu1, reconstructed from metagenome sequences, has been deposited in the Integrated Microbial Genomes (IMG) database (IMG Taxon ID: 2556793040) and the GenBank database (LQAA00000000).

### Classification

*Bacteria* » *Pseudomonadota* » *Deltaproteobacteria* » “Adiutricales” » “Adiutricaceae” » *Candidatus* *Adiutrix* » *Candidatus* *Adiutrix* intracellularis

### References

Effective publication: Ikeda-Ohtsubo et al., 2016 [1]

### Registry URL

<https://seqco.de/i:424>

## References

1. Ikeda-Ohtsubo et al. (2016). ‘*Candidatus* *Adiutrix* intracellularis’, an endosymbiont of termite gut flagellates, is the first representative of a deep-branching clade of *Deltaproteobacteria* and a putative homoacetogen. *Environmental Microbiology*. DOI:10.1111/1462-2920.13234