Register list for 3 new names including Ulvibacter alkanivorans sp. nov.

Submitted by Nguyen, Huong

Table 1: Complete list of names proposed in the current register list.

Proposed Taxon	Etymology	Description	Parent Taxon	Type	Registry URL
Species *Colwellia aromaticivorans*	[a.ro.ma.ti.ci.vo'rans] **L. masc. adj.** *aromaticus*, aromatic; **L. pres. part.** *vorans*, devouring; **N.L. fem. part. adj.** *aromaticivorans*, devouring aromatic compounds	Campeão et al. (2019): *In silico*-predicted phenotypes indicate that the novel species hydrolyzes chitin, utilizes propionate and dl-lactate, and does not utilize d-fructose, l-arabinose, glycerol, citrate, or l-malate. The G+C content is 38%.	*Colwellia*	NCBI Assembly: GCA_003333305.1 Ts	seqco.de/i:272
Species *Halocynthiibacter alkanivorans*	[al.ka.ni.vo'rans] **N.L. neut. n.** *alkanum*, alkane; **L. pres. part.** *vorans*, devouring; **N.L. masc. part. adj.** *alkanivorans*, devouring alkanes	Campeão et al. (2019): *In silico*-predicted phenotypes indicate that the novel species produces acid from d-ribose, sucrose, and glycerol. Acetate is utilized, nitrate is reduced, and indole is produced. The G+C content is 59%.	*Halocynthiibacter*	NCBI Assembly: GCA_003333265.1 Ts	seqco.de/i:273
Species *Ulvibacter alkanivorans*	[al.ka.ni.vo'rans] **N.L. neut. n.** *alkanum*, alkane; **L. pres. part.** *vorans*, devouring; **N.L. masc. part. adj.** *alkanivorans*, devouring alkanes	Campeão et al. (2019): *In silico*-predicted phenotypes indicate that the novel species utilizes dl-lactate and does not hydrolyze starch or carboxymethyl cellulose (CM-cellulose). It is negative for the utilization of d-mannitol, cellobiose, d-fructose, l-fucose, d-galactose, myo-inositol, d-sorbitol, sucrose, trehalose, and propionic acid. The G+C content is 40%.	*Ulvibacter*	NCBI Assembly: GCA_003333325.1 Ts	seqco.de/i:274