Genus Anaerocaecibacter

Etymology

[An.ae.ro.cae.ci.bac.ter.] N.L. masc. n. Anaerocaecibacter, an anaerobic rod from caecum

Nomenclatural type

Species Anaerocaecibacter muris[™]

Description

The closest relatives based on 16S rRNA gene similarity are *Xylanivirga thermophila* (83.57%), *Christensenella hongkongensis* (83.57%), *Caldicoprobacter guelmensis* (83.41%), and *Caldicoprobacter faecalis* (83.29%). POCP values to all close relatives were below 30% and GTDB-Tk placement assigned the type species to an unknown genus within 'f_CAG-552'. Separation from the other proposed species *Pumilibacter muris* and *Pumilibacter intestinalis* within the propose novel family *Pumilibacteraceae* (see protologue below) was confirmed by phylogenomic placement, which showed they were distinct from each other, and by the POCP value of 45.9% between the type species of each genus. The type species is *Anaerocaecibacter muris*.

Classification

Bacteria » Bacillota » Clostridia » Eubacteriales » Pumilibacteraceae » Anaerocaecibacter

References

Effective publication: Afrizal et al., 2022 [1]

Registry URL

https://seqco.de/i:23523

References

1. Afrizal et al. (2022). Enhanced cultured diversity of the mouse gut microbiota enables custom-made synthetic communities. *Cell Host & amp; Microbe*. DOI:10.1016/j.chom.2022.09.011