

Species *Methanocatella woesei*

Etymology

[woe'se.i] N.L. gen. masc. n. *woesei*, of Woese, named in honor of Carl R. Woese for his pioneering contributions to the understanding of the phylogeny of methanogens and other microorganisms

Nomenclatural type

[NCBI Assembly: GCF_003111605.1](#)^{Ts}

Reference Strain

[DSM 11979](#)

Description

Coccobacillus with slightly tapered or rounded ends, about 0.6 µm in width and 1.0 µm in length, occurring in pairs or short chains. Gram-positive reaction. Cell walls are composed of pseudomurein. Cells are resistant to lysis by SDS. Optimum temperature 37 °C. Optimum pH 7 (as in [Miller, Lin, 2002](#)). The G+C content of the type genome is 29.9 mol%, and the genome size is 1.54 Mbp.

Classification

Archaea » *Methanobacteriota* » *Methanobacteria* » *Methanobacteriales* » *Methanobacteriaceae* » *Methanocatella* » *Methanocatella woesei*

References

Effective publication: Protasov et al., 2023 [1]

Registry URL

<https://seqco.de/i:32438>

References

1. Protasov et al. (2023). Diversity and taxonomic revision of methanogens and other archaea in the intestinal tract of terrestrial arthropods. *Frontiers in Microbiology*. DOI:10.3389/fmicb.2023.1281628