

## Species *Methanolapillus ohkumae*

### Etymology

[oh.ku'mae] N.L. gen. masc. n. *ohkumae*, of Ohkuma, named after Moriya Ohkuma in recognition of his important contributions to arthropod gut microbiology

### Nomenclatural type

[NCBI Assembly: GCA\\_032594355.1<sup>Ts</sup>](#)

### Reference Strain

[DSM 114424 = JCM 39382](#)

### Description

Irregular cocci cells 1–2 µm in diameter, non-motile. Use methanol and methylamines as methanogenesis substrates together with hydrogen. Requires acetate, formate, and coenzyme M; yeast extract is stimulatory. The optimum temperature is 37 °C. The G+C content of the type genome is 41.0%, and the estimated genome size is 1.84 Mbp.

### Classification

Archaea » *Methanobacteriota* » “Methanomicrobia” » *Methanosarcinales* » *Methanosarcinaceae* » *Methanolapillus* » *Methanolapillus ohkumae*

### References

Effective publication: Protasov et al., 2023 [1]

### Registry URL

<https://seqco.de/i:32603>

## References

1. Protasov et al. (2023). Diversity and taxonomic revision of methanogens and other archaea in the intestinal tract of terrestrial arthropods. *Frontiers in Microbiology*. DOI:[10.3389/fmicb.2023.1281628](https://doi.org/10.3389/fmicb.2023.1281628)