

Species *Salinivivens marinus*^{Ts}

Etymology

[ma.ri'inus] L. masc. adj. *marinus*, of the sea

Nomenclatural type

[NCBI Assembly: GCA_038142605.1](#)^{Ts}

Description

Inferred to break down a wide range of organic carbon substrates (Chitin, Pectin, Cellulose and Starch). Capable of performing aerobic anoxygenic photosynthesis (pufLM), and thiosulfate oxidation (soxABCDXYZ), likely performing osmoregulation using trk1 (K⁺ transporter) and proXV (Glycine betaine) and inferred to be motile (fliGMN, motAB gene). Also carries genes for phosphorus uptake and regulation (pho, pst). Based on the genome reporting standards for MAGs, the estimated completeness 94.41%, contamination 0.3%, and the presence of the 5S (108 bp), 16S (1,457 bp), and 23S (2,718bp) rRNA gene and 32 tRNAs. Type genome is defined as “high-quality” draft MAG, with genome size of 2.6 Mbps

Classification

Bacteria » *Pseudomonadota* » *Alphaproteobacteria* » *Rhodobacterales* » *Paracoccaceae* » *Salinivivens* » *Salinivivens marinus*^{Ts}

References

Effective publication: Prabhu et al., 2024 [1]

Registry URL

<https://seqco.de/i:46736>

References

1. Prabhu et al. (2024). Machine learning and metagenomics identifies uncharacterized taxa inferred to drive biogeochemical cycles in a subtropical hypereutrophic estuary. *ISME Communications*.
[DOI:10.1093/ismeco/ycae067](https://doi.org/10.1093/ismeco/ycae067)