

Species *Hepatoplasma crinochetorum*^{Ts}

Etymology

[cri.no.che.to'rum] N.L. gen. pl. n. *crinochetorum*, of isopods (Crinocheta)

Nomenclatural type

[NCBI Assembly: GCA_000582535.1](#)^{Ts}

Description

Wang et al., 2004: spherical cells (diameter, 0.5 to 0.8 µm); no cell wall and outer membrane; forms stalk-like cytoplasmic appendages at one or two cell poles that attach to host epithelia or are involved in budding. Basis of assignment: 16S ribosomal DNA sequences (accession numbers AY500249 and AY500250), hybridization with 16S rRNA-targeted oligonucleotide probe (5'-GTG AAA AAT TCC CTA CTG CTG-3'). Symbiont of *P. scaber* and other terrestrial isopods belonging to the taxon Crinocheta, located exclusively in the lumen of the hepatopancreas; so far uncultivated.

Classification

Bacteria » *Mycoplasmata* » *Mollicutes* » *Mycoplasmatales* » *Hepatoplasmataceae* » *Hepatoplasma* » *Hepatoplasma crinochetorum*^{Ts}

References

Effective publication: Wang et al., 2004 [1]

Registry URL

<https://seqco.de/i:83>

References

1. Wang et al. (2004). " *Candidatus* *Hepatoplasma crinochetorum*," a New, Stalk-Forming Lineage of *Mollicutes* Colonizing the Midgut Glands of a Terrestrial Isopod. *Applied and Environmental Microbiology*. [DOI:10.1128/aem.70.10.6166-6172.2004](https://doi.org/10.1128/aem.70.10.6166-6172.2004)